(19) World Intellectual Property Organization International Bureau





(43) International Publication Date 7 February 2002 (07.02.2002)

PCT

(10) International Publication Number WO 02/10181 A1

(51) International Patent Classification⁷: C07H 17/08, A61K 31/7052, A61P 31/04

(21) International Application Number: PCT/IB01/01327

(22) International Filing Date: 25 July 2001 (25.07.2001)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data: P-200001934

31 July 2000 (31.07.2000) ES

(71) Applicant (for all designated States except US): QUIM-ICA SINTETICA, S.A. [ES/ES]; C. de l'Acer, 24, E-08004 Barcelona (ES).

(72) Inventors; and

- (75) Inventors/Applicants (for US only): BAÑON PARDO, Gabriel [ES/ES]; c/ Río Segura, 20, E-28803 Alcalá de Henares (ES). GONZALES HERNANDEZ, Pedro [ES/ES]; Avda. Reyes Católicos, 3A, 3°C, E-28802 Alcalá de Henares (ES). LOPEZ GARCIA, Julia [ES/ES]; c/ Andrés Mellado, 13, 1°C, E-28015 Madrid (ES). RIAÑO RIBOTA, Ana María [ES/ES]; Avda. del Mediterráneo, 17, E-28007 Madrid (ES). GELPI VINTRO, José Maria [ES/ES]; Ronda General Mitre, 33, 4°, E-08017 Barcelona (ES). FAU DE CASA-JUANA MUÑOZ, Miquel [ES/ES]; c/ Guaimaral, 4, 2° derecha, E-28033 Madrid (ES).
- (74) Agents: PONTI SALES, Adelaida et al.; C. Consell de Cent, 322, E-08007 Barcelona (ES).
- (81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

- as to the identity of the inventor (Rule 4.17(i)) for the following designations AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG)
- as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii)) for the following designations AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG)
- of inventorship (Rule 4.17(iv)) for US only

Published:

- with international search report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: AZITHROMOCIN MONOHYDRATE WITH LOWER HYGROSCOPICITY, PREPARATION METHOD AND PHARMACEUTICAL COMPOSITIONS WHICH INCLUDE IT

(57) Abstract: The improved form of azitromycin monohydrate of lower hydroscopicity and greater density and hardness that presents values of: apparent filling density not lower than 0.35 g/ml; and apparent knocked density not lower than 0.50 g/ml. The method for preparation of an improved form of azitromycin monohydrate is characterised in that the hydroscopic azitromycin monohydrate is subjected to a procession of compaction by pressure. The invention also refers to a pharmaceutical composition that includes a therapeutically effective quantity of improved form of azitromycin monohydrate in association with at least one pharmaceutically acceptable suitable inert diluent.



A1

1

AZITHROMOCIN MONOHYDRATE WITH LOWER HYGROSCOPICITY, PREPARATION METHOD AND PHARMACEUICAL COMPOSITIONS WHICH INCLUD IT

5

Field of the invention

This invention relates to a new improved form of azitromycin (9-deoxo-9a-aza-9a-methyl-9a-homoerythromycin 10 A) monohydrate, the monohydrated form of azitromycin of lower hygroscopicity, the method for making it and the pharmaceutical compositions which include it.

Background of the invention

15

Azitromycin or 9-deoxo-9a-aza-9a-methyl-9ahomoerythromycin A is a broad-spectrum antibacterial agent that was disclosed and patented by Sour Pliva in Yugoslav priority application YU 000592 of 06.03.81, which priority 20 was invoked in the equivalent American patent US 4517359.

2

These patents describe the preparation of the product by means of an Eschweiler-Clarke methylation reaction of 9-deoxo-9a-aza-9a-methyl-9a-homoerythromycin A.

5 Two crystalline forms of azitromycin are known: azitromycin monohydrate and azitromycin dihydrate.

dihydrated form of azitromycin is obtained by The crystallisation of the product in a mixture of and an aliphatic hydrocarbon in the 10 tetrahydrofuran (EP 298.650) or by means of presence of water recrystallisation of the product of aketone and water (J. Chem. Research (M), 1988, 1239-1261; J. Chem. Research (S), 1988, 152-153).

15

Furthermore, the azitromycin monohydrate is prepared by crystallisation of the product from mixtures of ethanol and water (EP 298.650) or aketone and water.

- 20 Both monohydrated and dihydrated forms are easily distinguishable by their X-ray diffraction spectrums, IR spectrums, differential calorimetry analysis (DSC) and thermogravimetric analysis (TGA).
- 25 In European patent EP 298.650 from Pfizer, Messrs Allen and Nepveux describe how when azitromycin is obtained according to the Eschweiler-Clark methylisation procedure of 9-deoxo-9a-aza-9a-methyl-9a-homoerythromycin A with formaldehyde and formic acid and the azitromycin resulting
- 30 from ethanol and water is recrystallised, the monohydrated form of azitromycin is obtained. This product is hygroscopic and can be used only with difficulty, since the monohydrate absorbs variable amounts of water, which makes it difficult to use for medicinal formulation.

WO 02/10181

3

The only method by which Messrs Allen and Nepveux overcame these difficulties and limitations of the monohydrated form of azitromycin was by developing a new form of azitromycin, the dihydrate of azitromycin, described and 5 claimed in EP 298.650.

Description of the invention

It is, therefore, an object of this invention to provide 10 an improved form of azitromycin monohydrate which overcomes the difficulties and limitations of the hygroscopic azitromycin monohydrate form previously known and which shows lower hygroscopicity.

- · 15 The term "an improved form of azitromycin monohydrate" can alternatively be denominated "azitromycin monohydrate of the invention". These terms are used interchangeably in this specification.
 - 20 The term "hygroscopic azitromycin monohydrate" refers to the azitromycin monohydrate previously known through EP298650. More specifically, it refers to the azitromycin monohydrate described on page 4, 'lines 25-48 of EP298650.
 - 25 This "hygroscopic azitromycin monohydrate" is the reference molecule of this invention. In consequence, when for example in the invention "an improved form of azitromycin monohydrate that presents lower hygroscopicity" is used, this "lower hygroscopicity"
 - 30 refers to lower hygroscopicity when compared with the hygroscopicity of the reference molecule "hygroscopic azitromycin monohydrate" reference molecule of EP298650. The same is the case when the statement "denser and stronger" is used in relation to the azitromycin of the

4

invention, that is, it is "denser" when compared with the reference molecule of EP298650.

Another object of this invention is to disclose a method 5 of preparation of an improved form of monohydrated azitromycin of lower hygroscopicity, which method provides the improved form of monohydrated azitromycin with a practically constant water content and in a reproducible manner.

10

Another object of this invention is to disclose pharmaceutical compositions which include an improved form of azitromycin monohydrate, the monohydrated azitromycin of lower hygroscopicity.

15

In accordance with one of the objects of this invention, a new form of azitromycin monohydrate has been surprisingly discovered, a form which overcomes the difficulties and limitations of the previously known form of azitromycin 20 monohydrate and which presents lower hygroscopicity, this improved form being a compound of lower hygroscopicity which has apparent density values that are especially suitable for its use as an active ingredient in the formulation of tablets and other medicinal forms.

25

The improved form of azitromycin monohydrate of lower hygroscopicity is characterised in that it is denser and stronger and possesses values of:

- apparent filling density not lower than 0.35 30 g/ml, preferably not lower than 0.40 g/ml, and more preferably still not lower than 0.45 g/ml, and still more preferably not lower than 0.50 g/ml and yet more preferably not lower than 0.53 g/ml; and - apparent knocked density not lower than 0.50 g/ml, preferably not lower than 0.55 g/ml, and more preferably still not lower than 0.60 g/ml, and still more preferably not lower than 0.65 g/ml and yet more 5 preferably not lower than 0.70 g/ml.

Preferably, the "apparent filling density" is measured by weighing out approximately 100 g of product measured with a precision of 0.1%. If it is not possible to use 100 g, 10 the quantity of sample and the volume of the cylinder and the conditions specified in the analysis can be altered. A quantity of sample is selected such that it has an apparent loose volume of 150 to 250 ml. The uncompacted powder is then transferred carefully to a dry graduated 15 cylinder of 250 ml (graduated cylinder weight: 220±44 g), by sliding the product gently. The apparent unsedimented volume occupied by the sample is then read, and it is approximated to the closest graduated unit: Volume (1) (expressed in ml), and the apparent filling density 20 calculated using the formula:

Apparent filling density = weight of sample (g)/V(1) ml. This procedure is that of USP24, Test 616.

Preferably, the apparent knocked density is measured in a Pharma Test apparatus (model PT-TD), with a knocking speed of 250±10 blows/minute and a knocking height of 3±0.1mm, securing the cylinder in the bracket of the equipment and knocking initially 500 times. The volume occupied by the 30 sample is noted and approximated to the nearest graduation of the graduated cylinder. This is the volume (2). The sample is then given a further 750 knocks and the knocked volume occupied by the sample measured: Volume (3). If the difference between the two volumes [V8·9-V(2)] is less 35 than 2%, V(3) is the final volume for calculating the

6

apparent knocked density. If the difference is greater than 2%, then knocking of the sample in increments of 1250 blows each time is continued until consecutive readings with a difference of less than 2% between them are obtained. The knocked density in g/ml is calculated using the following formula:

Apparent knocked density = weight of sample (g)/V(3) ml. This procedure is that of USP24, Test 616.

10

The technical characteristic of the density of the azitromycin monohydrate of the invention provides the effect of lower hygroscopicity. The object of this invention, as mentioned above, can therefore be expressed 15 alternatively as: a azitromycin monohydrate which is characterised in that is possesses values of:

- apparent filling density not lower than those mentioned previously; and
- apparent knocked density not lower than those mentioned previously.

The improved form of azitromycin monohydrate of lower hygroscopicity is further characterised in that it presents an initial water content of between 3.2% and 25 3.6%, a water gain of between 0.16 and 0.23% at 24 hours and between 0.3 and 0.51% at four days, finally stabilising at a water content of 3.6 to 3.9%, under ambient humidity conditions of between 20 and 40% and at temperatures in the range 20-30°C.

30

The azitromycin monohydrate presents apparent filling density values not exceeding 0.29 g/ml and apparent knocked density values not exceeding 0.44 g/ml.

7

Moreover, if the hygroscopic azitromycin monohydrate is exposed to the atmosphere and maintained under mean ambient conditions of 20-30°C of temperature and 20-40% of ambient relative humidity, it presents a water gain of 5 1.2% at 24 hours and of 1.3% at four days, finally stabilising at a water content of 4.6%.

The X-ray diffraction spectrum of the improved form of azitromycin monohydrate of lower hygroscopicity shows that 10 the product presents a lower crystalline content when compared with that of the hygroscopic azitromycin monohydrate, which can have a positive effect in improvements in its solubility and bioavailability.

- 15 Furthermore, the values for melting point, DSC and IR spectrum of the new improved form of azitromycin monohydrate of lower hygroscopicity are identical to those of the hygroscopic azitromycin monohydrate.
- 20 In accordance with another aspect of this invention, a method is disclosed for preparation of an improved form of azitromycin monohydrate, the azitromycin monohydrate of lower hygroscopicity, which method gives rise surprisingly to an improved form of azitromycin monohydrate with a 25 practically constant water content and in a reproducible manner.

The hygroscopic azitromycin monohydrate is obtained by means of an Eschweiler-Clarke reaction of the 9-deoxo-9a-30 aza-9a-methyl-9a-homoerythromycin A with formaldehyde and formic acid, the resulting product being recrystallised in mixtures of ethanol and water or of acetone and water to provide the hygroscopic monohydrate form.

8

The method for preparing an improved form of azitromycin monohydrate, the azitromycin monohydrate of lower hygroscopicity, is characterised in that the hygroscopic azitromycin monohydrate is subjected to a procession of 5 compaction by pressure.

The hygroscopic azitromycin monohydrate can be obtained via the reaction based on Eschweiler-Clark described above. However, it is possible to obtain it by different 10 protocols in accordance with the general state of the art on the subject. The characteristic feature of the method of the invention is that the hygroscopic azitromycin monohydrate is submitted to a process of compaction by pressure. The specific method used for obtaining the 15 hygroscopic azitromycin monohydrate is not essential in nature, although the preferred procedure is that mentioned previously.

The compaction pressure can range between 50 and 200 20 kg/cm2, but is preferably between 100 and 200 kg/cm2.

The speed of rotation of the rollers has to be between 5 and 15 rpm, and is preferably between 9 and 15 rpm.

25 The mesh size has to be between 0.5 and 2 mm, preferably with an opening size of between 1 and 2 mm.

Any type of compactor can be used, though preferably a roller compactor.

30

In accordance with the method object of this invention, it has been found that when the hygroscopic azitromycin monohydrate is subjected to a process of compaction by pressure resulting in a product of greater apparent 35 density, this new product surprisingly presents the

9

advantageous property of possessing a lower tendency to take up water contained in the atmosphere, the characteristic of lower hygroscopicity.

5 This new improved form of azitromycin monohydrate of lower hygroscopicity remains stable during and after the process of compaction by pressure, without there taking place any increase in the level of impurities in the resulting product.

10

In accordance with the method of this invention, it has surprisingly proved possible to obtain a product which, if exposed to the atmosphere and kept at mean ambient conditions of 20-30°C of temperature and 20-40% of ambient 15 relative humidity, presents a water gain of 0.23% at 24 hours and of 0.513% at four days, finally stabilising at a water content value of 3.9%, while the uncompacted hygroscopic azitromycin monohydrate presents a water gain of 1.2% and 1.3%, respectively, at the end of said periods 20 of time, finally stabilising at a water content of 4.6%.

The efficacy of the method in this respect was confirmed by the fact when the same product is submitted to a double operation of compaction by means of pressure, the water 25 gain after 24 hours and 4 days is 0.16% and 0.30%, respectively, leaving the product finally stabilised at a water content of 3.6%.

Another object of this invention relates to the 30 pharmaceutical composition that include a therapeutically effective quantity of azitromycin monohydrate of lower hygroscopicity in association with at least one pharmaceutically acceptable suitable inert diluent.

10

The pharmaceutical composition in accordance with this invention can be prepared by combining the azitromycin monohydrate of lower hygroscopicity with at least one pharmaceutically acceptable suitable inert diluent and 5 administering it orally and parenterally.

This invention is illustrated by means of the following examples. It should nevertheless be understood that this invention is not limited to the specific examples in them.

10

Drawings

Figure 1 shows the X-ray spectrum of hygroscopic azitromycin monohydrate.

15

Figure 2 shows the X-ray diffractogram of the improved form of azitromycin monohydrate.

Figure 3 shows the IR spectrum of the hygroscopic 20 azitromycin monohydrate.

Figure 4 shows the IR spectrum of the improved form of azitromycin monohydrate of lower hygroscopicity.

25 Examples

Example 1 to 5

20 Kg of hygroscopic azitromycin monohydrate having 30 filling density values of 0.29 g/ml and knocked density of 0.44 g/ml were compacted in a roller compactor of the Bonal trademark, model BC-175/75V fitted with screen with 1 mm mesh size, and the compaction pressure and roller rotation speed parameters were set as indicated in Table

A. The density results obtained are shown in the same table.

Table A: Compaction Conditions

5

Example	Compaction	Roller	Filling	Knocked
No.	Pressure	rotation	Density	Density
	(kg/cm2)	speed	(g/ml)	(g/ml)
		(rpm)		
1	75	11.1	0.49	0.61
2	100	11.1	0.51	0.62
3	130	10	0.52	0.64
4	150	10	0.53	0.70
5 (*)	150	10	0.55	0.74

(*) New product compaction value obtained by combining examples 1 and 2.

10

Hygroscopicity results

Table B below shows the comparative hygroscopicity results obtained with the hygroscopic azitromycin monohydrate and 15 the improved form of azitromycin of lower hygroscopicity. Said table shows the changes in the water content of the product over time, measured by the Karl Fischer method, when exposed to the following ambient conditions:

Temperature between: 20-30°C

20 Relative Humidity between: 20-40%

12

Table B: Hygroscopicity

Time	Hygroscopic azitromycin monohydrate	Product Example 4	Product Example 5	
	(% H ₂ O)	(% H ₂ O)	(% H ₂ O)	
0	3.28	3.39	3.33	
24 hours	4.49	3.62	3.49	
2 days	4.52	3.70	3.53	
4 days	4.59	3.90	3.63	
15 days	4.6	3.9	3.6	

5

Stability results

Table C below shows that the quality of the product remained unchanged after it had been submitted to the 10 compaction process.

Table C: Stability

Parameter	Hygroscopic azitromycin monohydrate	Product Example 4
Specific rotation (s.s.s.)	-48.8°	-47.5°
pH (0.2% solution in methanol/water, 1:1)	9.98	10.1
Purity (%)	98.0	98.2

13

Results of the X-ray diffractograms

The X-ray diffractograms were obtained by means of a 5 Philips Analytical X-Ray B.V. diffractometer of type PW 1710, operating at 40 kV and 40 mA and using a copper radiation of $k\alpha 1=1-54056A$ and $k\alpha 2=1.54439A$. This equipment has a graphite monochromator with automatic slit, and the crystalline powder method was used.

10

Figure 1 corresponds to the X-ray spectrum of hygroscopic azitromycin monohydrate, while Figure 2 corresponds to the X-ray diffractogram of the improved form of azitromycin monohydrate.

15

A comparison of them shows a reduction of crystallinity in the improved form of azitromycin monohydrate.

20 Infrared spectrum results

Attached below are the IR spectrum results of hygroscopic azitromycin monohydrate and the improved form of azitromycin monohydrate, recorded on a Perkin Elmer 25 spectrometer, model 1600 Series FTIR.

Figure 3 shows the IR spectrum of the hygroscopic azitromycin monohydrate, while Figure 4 shows the IR spectrum of the improved form of azitromycin monohydrate 30 of lower hygroscopicity.

14

CLAIMS

- Improved form of azitromycin monohydrate of lower hydroscopicity and greater density and hardness that 5 presents values of:
 - apparent filling density not lower than 0.35 g/ml; and
 - apparent knocked density not lower than 0.50 g/ml.

10

- 2. Improved form of azitromycin monohydrate as claimed in Claim 1, which further presents an initial water content of between 3.2% and 3.6%, a water gain of between 0.16 and 0.23% at 24 hours and between 0.30 and 15 0.51% at four days, finally stabilising at a water content of 3.6 to 3.9%, under ambient humidity conditions of between 20 and 40% and at temperatures in the range 20-30°C.
- 3. Improved form of azitromycin monohydrate as claimed in Claim 1, in which the apparent filling density is not less than 0.40 g/ml and apparent knocked density not less than 0.55 g/ml.
- 4. Improved form of azitromycin monohydrate as claimed in Claim 1, in which the apparent filling density is not less than 0.45 g/ml and apparent knocked density not less than 0.60 g/ml.
- 5. Improved form of azitromycin monohydrate as claimed in Claim 1, in which the apparent filling density is not less than 0.50 g/ml and apparent knocked density not less than 0.65 g/ml.

15

6. Improved form of azitromycin monohydrate as claimed in Claim 1, in which the apparent filling density is not less than 0.53 g/ml and apparent knocked density not less than 0.70 g/ml.

5


- 7. Method for preparation of an improved form of azitromycin monohydrate as claimed in any of the previous claims, characterised in that the hygroscopic azitromycin monohydrate is subjected to a procession of compaction by 10 pressure.
- 8. Method as claimed in Claim 7, characterised in that the compaction pressure is furthermore ranges between 50 and 200 kg/cm 2 , the speed of rotation of the rollers is 15 between 5 and 15 rpm and the mesh size is between 0.5 and 2 mm.
- 9. Method as claimed in Claim 8, characterised in that the compaction pressure is $150~kg/cm^2$, the speed of 20 rotation of the rollers is 10~rpm and the mesh size is 1~mm.
 - 10. Product that can be obtained according to any of method claims 7, 8 or 9.

25

- 11. Pharmaceutical composition that includes a therapeutically effective quantity of improved form of azitromycin monohydrate as claimed in any of Claims 1 to 6 in association with at least one pharmaceutically 30 acceptable suitable inert diluent.
- 12. Method for preparation of the pharmaceutical composition claimed in Claim 11, characterised in that it combines the azitromycin monohydrate of lower 35 hygroscopicity with at least one pharmaceutically

16

acceptable suitable inert diluent, administered orally and parenterally.








FIG. 3







INTERNATIONAL SEARCH REPORT

Internati Application No PCT/IB 01/01327

A. CLASSIF	CATION OF SUBJECT MATTER C07H17/08 A61K31/7052 A61P31/04		
166 /	CU/H17/U8 AG1K31//U52 AG1F31/U4		
According to	International Palant Classification (IDC) and both patients described	lon and IDC	
B. FIELDS	International Patent Classification (IPC) or to both national classification	ori and IPO	
	cumentation searched (classification system followed by classification	n symbols)	
IPC 7	CO7H A61K A61P		
Documentati	on searched other than minimum documentation to the extent that su	ch documents are included in the fields per	arched
,	Or searched when than minimum documentation to the exicit mar so	on documento die molidada in tila liena des	a died
Electronic da	ata base consulted during the international search (name of data base	e and, where practical search terms used)	
	ternal, WPI Data, CHEM ABS Data	, , ,	
	, 2004, 0		ł
		•	ļ
C. DOCUME	ENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant	vant passages	Relevant to claim No.
	ED 0 000 CEO 4 (PETZED)		
A	EP 0 298 650 A (PFIZER) 11 January 1989 (1989-01-11)		1
	cited in the application		
	the whole document		
P,X	WO 01 00640 A (LUDESCHER JOHANNES		1,10-12
	RAFAEL (ES); BIOCHEMIE SA (ES); Di 4 January 2001 (2001-01-04)	IAGO J)	
	abstract		
}	claims		
·	din ani un vinann		
			İ
ļ			
Ì			
Furt	her documents are listed in the continuation of box C.	Patent family members are listed i	n annex.
° Special ca	ategories of cited documents:	"T" later document published after the inter or priority date and not in conflict with	
A docum	ent defining the general state of the art which is not dered to be of particular relevance	cited to understand the principle or the invention	
E earlier	document but published on or after the international date	"X" document of particular relevance; the cl cannot be considered novel or cannot	laimed invention be considered to
which	ent which may throw doubts on priority claim(s) or is cited to establish the publication date of another	involve an inventive step when the doc "Y" document of particular relevance; the c	cument is taken alone
O, docum	on or other special reason (as specified) ent referring to an oral disclosure, use, exhibition or	cannot be considered to involve an inv document is combined with one or mo	ventive step when the re other such docu-
P docum	means ent published prior to the international filing date but	ments, such combination being obviou in the art.	
	han the priority date claimed actual completion of the international search	*&" document member of the same patent in Date of mailing of the international sea	
5 September 2001		19/09/2001	
Name and	mailing address of the ISA European Patent Office, P.B. 5818 Patentiaan 2	Authorized officer	
	NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Tx. 31 651 epo nt,	do Nooy A	
	Fax: (+31-70) 340-3016	de Nooy, A	

INTERNATIONAL SEARCH REPORT

Information on patent family members

Internati spplication No
PCT/IB 01/01327

					01/0132/
Patent document cited in search report		Publication Patent family Publication date member(s) Publication date			
EP 0298650	A 11-01-	1989 WO	89005		26-01-1989
		AP		44 A	27-07-1989
		AT	724	46 T	15-02-1992
		AU	6045	53 B	20-12-1990
		AU	18839	88 A	12-01-1989
		BA	982	13 B	02-08-1999
		BG	473	48 A	15-06-1990
		CA	13148	76 A	23-03-1993
		CN	10304	22 A,B	18-01-1989
		CS	88048	96 A	14-03-1990
		CY		76 A	20-10-1995
		DD	2717		13-09-1989
		DE	38682	96 A	19-03-1992
		DK	3806	88 A	10-01-1989
		ES	20387	56 T	01-08-1993
		FI		87 A,B,	08-01-1990
		GR			16-03-1993
		HK		94 A	25-11-1994
		HU	95007		28-11-1995
		IE		54 B	29-06-1994
		IL		79 A	15-11-1992
		IN		379 A	29-06-1991
		JP	10380		08-02-1989
		JP	19035		08-02-1995
		JP	60313		27-04-1994
		KR			25-08-1990
		LV		24 A	20-04-1995
		MX		213 A	01-05-1993
		NZ		38 A	26-02-1990
		OA.		43 A	31-03-1989
		PT		33 A,B	30-06-1989
		RO			30-10-1993
		SG		794 G	14-10-1994
		SI			31-12-1996
		RU			10-09-1996
		US			31-07-2001
		YU		88 A	28-02-1990
		ZA 	88049	925 A	28-02-1990
WO 0100640	A 04-01	-2001 AU	58204	100 A	31-01-2001

		•
		•